

# *Il2rb* Cas9-CKO Strategy

**Designer:** Jing Jin

**Reviewer:** Yang Zeng

**Design Date:** 2018-6-22

# Project Overview

**Project Name**

***Il2rb***

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Il2rb* gene. The schematic diagram is as follows:



# Technical routes

- The *Il2rb* gene has 2 transcripts. According to the structure of *Il2rb* gene, exon3-exon9 of *Il2rb-201* (ENSMUST00000089398.8) transcript is recommended as the knockout region. The region contains 824bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Il2rb* gene. The brief process is as follows: sgRNA was transcribed in vitro, donor vector was constructed. Cas9, sgRNA and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice was knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Homozygotes for a targeted null mutation exhibit spontaneous activation of T cells and differentiation of B cells, elevated immunoglobulins including autoantibodies causing hemolytic anemia, granulocytopenia, and death after 3 months of age.
- The *Il2rb* gene is located on the Chr15. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

# Gene information (NCBI)

## Il2rb interleukin 2 receptor, beta chain [Mus musculus (house mouse)]

Gene ID: 16185, updated on 19-Mar-2019

### Summary



<b>Official Symbol</b>	Il2rb provided by <a href="#">MGI</a>
<b>Official Full Name</b>	interleukin 2 receptor, beta chain provided by <a href="#">MGI</a>
<b>Primary source</b>	<a href="#">MGI:MGI:96550</a>
<b>See related</b>	<a href="#">Ensembl:ENSMUSG00000068227</a>
<b>Gene type</b>	protein coding
<b>RefSeq status</b>	REVIEWED
<b>Organism</b>	<a href="#">Mus musculus</a>
<b>Lineage</b>	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
<b>Also known as</b>	CD122, IL-15Rbeta, IL15Rbeta, IL-2/15Rbeta, IL-2Rbeta, p70
<b>Summary</b>	The interleukin 2 receptor is composed of alpha and beta subunits. The beta subunit encoded by this gene is very homologous to the human beta subunit and also shows structural similarity to other cytokine receptors. [provided by RefSeq, Jul 2008]
<b>Expression</b>	Biased expression in spleen adult (RPKM 30.6), mammary gland adult (RPKM 14.0) and 8 other tissues <a href="#">See more</a>
<b>Orthologs</b>	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 2 transcripts, all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
<b>Il2rb-201</b>	<a href="#">ENSMUST00000089398.8</a>	4194	<a href="#">539aa</a>	Protein coding	<a href="#">CCDS27616</a>	<a href="#">P16297</a>	TSL:1 GENCODE basic APPRIS P1
<b>Il2rb-202</b>	<a href="#">ENSMUST00000163494.2</a>	1895	<a href="#">539aa</a>	Protein coding	<a href="#">CCDS27616</a>	<a href="#">P16297</a>	TSL:1 GENCODE basic APPRIS P1

The strategy is based on the design of *Il2rb-201* transcript, The transcription is shown below



# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Homozygotes for a targeted null mutation exhibit spontaneous activation of T cells and differentiation of B cells, elevated immunoglobulins including autoantibodies causing hemolytic anemia, granulocytopenia, and death after 3 months of age.

If you have any questions, you are welcome to inquire.

Tel: 025-5864 1534

