

Sdc4 Cas9-KO Strategy

Designer: Yanhua Shen

Project Overview



Project Name

Sdc4

Project type

Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Sdc4* gene. The schematic diagram is as follows:



- The *Sdc4* gene has 2 transcripts. According to the structure of *Sdc4* gene, exon2 of *Sdc4-201* (ENSMUST00000017153.3) transcript is recommended as the knockout region. The region contains 127bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Sdc4* gene. The brief process is as follows: CRISPR/Cas9 system v

- According to the existing MGI data, Mice homozygous or heterozygous for a knock-out allele show delayed wound healing and impaired angiogenesis. Homozygotes for a different knock-out allele exhibit degenerated fetal vessels in the placental labyrinth, abnormal cell adhesion, and high susceptibility to induced renal and hepatic injury.
- The *Sdc4* gene is located on the Chr2. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)

Sdc4 syndecan 4 [Mus musculus (house mouse)]

Gene ID: 20971, updated on 25-Mar-2019

Summary




Official Symbol	Sdc4 provided by MGI
Official Full Name	syndecan 4 provided by MGI
Primary source	MGI:MGI:1349164
See related	Ensembl:ENSMUSG00000017009
Gene type	protein coding
RefSeq status	PROVISIONAL
Organism	Mus musculus
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as	AA959608, AW108331, S4, Synd4, ryudocan, syndecan-4
Expression	Broad expression in kidney adult (RPKM 79.7), liver adult (RPKM 72.9) and 24 other tissues See more
Orthologs	human all

Transcript information (Ensembl)


The gene has 2 transcripts, all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Sdc4-201	ENSMUST00000017153.3	2458	198aa	Protein coding	CCDS17036	Q35988 Q3U5S6	TSL:1 GENCODE basic APPRIS P1
Sdc4-202	ENSMUST00000142909.1	797	No protein	Processed transcript	-	-	TSL:3


The strategy is based on the design of *Sdc4-201* transcript, The transcription is shown below




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Mice homozygous or heterozygous for a knock-out allele show delayed wound healing and impaired angiogenesis. Homozygotes for a different knock-out allele exhibit degenerated fetal vessels in the placental labyrinth, abnormal cell adhesion, and high susceptibility to induced renal and hepatic injury.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

