


Genotyping Report

Strain ID	T052837	Strain Type	KO(Cas9)	Genetic Background	C57BL/6JGpt
Designer	Zifan Lin	Gene Name	<i>Il1rl2</i>		


1. Strategy of Genotyping



2. Primer Information

PCR No.	Primer No.	Sequence	Band Size
PCR①	T052837-F1	CCGAAGATGGCTAATGGTTTTGG	WT:2623bp KO: 404bp
	T052837-R1	CTACAAGACATTCGTGGAACCGTG	
PCR②	T052837-F2	TAGTTCAAGAGGTTTCATGGTGGCC	WT:382bp KO: 0bp
	T052837-R2	CTGGTGAATTTACGGGACTCCC	

3. Gel Image



- ① Control (WT) : It is an important reference mark for whether the PCR reaction is successful and whether the product band position and size meet the theoretical requirements.
- ② Control (B) : PCR amplification was performed without template in the PCR reagent to monitor whether the reagent was contaminated.

4. PCR Condition

(Generally recommend to use Vazyme P222; If the sequences contain special structures such as GC% \geq 60% or GC% \leq 40%, recommend to use Vazyme P515.)

PCR Reaction Component			
Seg.	Reaction Component		Volume (μ l)
1	2 \times Rapid Taq Master Mix(Vazyme P222) or 2 \times Phanta Max Master Mix (Vazyme P515)		12.5
2	ddH ₂ O		9.5
3	Primer A(10pmol/ μ l)		1
4	Primer B(10pmol/ μ l)		1
5	Template(20~80ng/ μ l)		1
PCR program I priority selection			
Seg.	Temp.	Time	Cycle
1	95 $^{\circ}$ C	5min	20 \times
2	98 $^{\circ}$ C	30s	
3	65 $^{\circ}$ C*(-0.5 $^{\circ}$ C/cycle)	30s	
4	72 $^{\circ}$ C	45s*	
5	98 $^{\circ}$ C	30s	15 \times
6	55 $^{\circ}$ C*	30s	
7	72 $^{\circ}$ C	45s*	
8	72 $^{\circ}$ C	5min	
9	10 $^{\circ}$ C	hold	
PCR program II the second choice			
Seg.	Temp.	Time	Cycle
1	95 $^{\circ}$ C	5min	35 \times
2	98 $^{\circ}$ C	30s	
3	58 $^{\circ}$ C*	30s	
4	72 $^{\circ}$ C	45s*	
5	72 $^{\circ}$ C	5min	
6	10 $^{\circ}$ C	hold	

Note*: Annealing temperature and extension time can be determined according to the actual amplification situation and amplification enzyme efficiency.

